

## New approximating results for data with errors in both variables

Nina B. Bogdanova<sup>1</sup> and Stefan T. Todorov.<sup>2</sup>  
*INRNE, BAS, 72 Tzarigradsko choussee, 1784 Sofia, Bulgaria*

### Abstract

We introduce new data from mineral water probe /Lenovo Bulgaria/, measured with errors in both variables. We apply our Orthonormal Polynomial Expansion Method (OPEM), based on Forsythe recurrence formula to describe the data in the new error corridor. The development of OPEM gives the approximating curves and their derivatives in optimal orthonormal and usual expansions including the errors in both variables with special criteria.

### Аннотация

Мы рассматриваем новые данные для пробы минеральной воды из источника в Леново (Болгария), измеренные с ошибками по обоим переменным. Для описания данных в новом коридоре ошибок применяется наш метод разложения по ортогональным полиномам, основанный на рекуррентной формуле Форсайта. Представлено дальнейшее развитие нашего численного метода, связанное с использованием критерия оптимального ортогонального разложения и оценкой на этой основе разложения обычного.

PACS: 02.30.-Mv; 02.60.-Ed ;02.60.-x;

## 1 Introduction

The water spectra method applies a drop taken from a water probe to measure the water's state spectrum. In the special experiment the drop is placed on a hostaphan folio- Fig. 1. During the whole process of evaporation of the drop, one measures at equal time intervals the drop contact angle with the folio. On independent axis one has the values of the contact angles within fixed angular intervals and on dependent axis the frequency of measurements of these angles.  $\phi(\theta)$  is the distribution of the number of measurements of contact angle  $\theta$  during the drop evaporation. One can change the function  $\phi(\theta)$  of the angle  $\theta$  to the function of energy variable  $F(E)$  using the following Antonov transformation [1, 2]:

$$E(\theta) = -\gamma(1 + \cos \theta)/I(1 + \cos \theta_0), F(E) = b\phi(\theta)/\sqrt{1 - (1 + bE)^2},$$

where  $b = I(1 + \cos(\theta_0))/\gamma$ . Here  $I = 5,03 \cdot 10^{18} \text{ m}^{-2}$  is the density of water molecules in the surface layer,  $\gamma$  is the surface tension,  $\theta_0$  - the initial contact angle. The so obtained graph after measurements by method in [1] is referred to as energy spectrum  $F(E)$  of the probe. The domain of  $F(E)$  contains the values of the Hydrogen bond energy.

The method of water spectra is sensitive to treatment by physical fields as e.g.  $\gamma$ -ray treatment of water filtration [1, 2] and to environmental changes of the ecosystem on and mechanical treatment and aeration on different water probes [3, 4]. In the present paper we approximate another natural water data taken from a water spring in Bulgaria near the village Lenovo. On Figure 2 we present new detailed information about given data and their errors on both variables of water probe.

<sup>1</sup>Email:nibogd@inrne.bas.bg

<sup>2</sup>Email:todorov\_st@yahoo.com




Fig. 1: Experimental setup




Fig. 2: Given data (B) with errors

## 2 Main problem definition

- To find the best approximation curve of measured water data on Fig.2 taking into account the errors in both variables;
- To extend our Orthonormal Polynomial Expansion Method (OPEM), according some criteria, to evaluate orthonormal description of given data. To find the best approximating curve with usual polynomials, received by orthonormal, according some criteria.

## 3 Numerical method–OPEM “total variance”

Let the  $\{E_i, F_i, i = 1, \dots, M\}$  are arbitrary pairs of monitoring data  $E$  and  $F$ , introduced in section 2. They are given with experimental errors in both variables-  $\sigma(F_i)$  and  $\sigma(E_i)$ . Consider the total uncertainty (total variance)  $S^2(E, F)$  [5, 6, 7], associated with  $(E, F)$

$$S_i^2 = \sigma^2(F_i) + \left(\frac{\partial F_i}{\partial E_i}\right)^2 \sigma^2(E_i), \quad (1)$$

according the ideas of Bevington (1977)[5], where his proposal is to combine the errors in both variables and assign them to dependent variable. One defines the errors corridor  $C(E, F)$ , which is the set of all intervals

$$[F(E) - S(E, F), F(E) + S(E, F)], \quad (2)$$

### 3.1 Orthonormal expansion criteria

The first criterion to be satisfied, is that the fitting curve should pass within the errors corridor  $C(E, F)$ . In the cases of errors only in  $F$ , (i.e.  $\sigma(E) = 0, \sigma(F) \neq (0)$ ) the errors corridor  $C(E, F)$  reduces to the known set of intervals

$$[F - \sigma(F), F + \sigma(F)], \quad (3)$$

for any  $F$ . The second criterion is, that the fitting curve  $F^{\text{appr}}(E_i)$  satisfies the expression

$$\chi^2 = \sum_{i=1}^M w_i [F^{\text{appr}}(E_i) - F(E_i)]^2 / (M - L), w_i = 1/S_i^2. \quad (4)$$

should be minimal (L-number of polynomials). The preference is given to the first criterion. When it is satisfied, the search of the minimal chi-squared stops. Some details of the calculation procedure are given in Forsythe's paper [7] and in our works [8, 9, 10].

Our procedure gives results for approximating function by two expansions : of orthogonal coefficients  $\{a_i\}$  with optimal degree  $L_a$  and usual ones  $\{c_i\}$  with optimal degree  $L_c$ :

$$F^{\text{appr}(m)}(E) = \sum_{i=0}^{L_a} a_i P_i^{(m)}(E) = \sum_{i=0}^{L_c} c_i E^i. \quad (5)$$

The orthogonal coefficients are evaluated by the given values  $F_i$ , weights and orthogonal polynomials:

$$a_i = \sum_{k=1}^M F_k w_k P_i^{(m)}(E_k). \quad (6)$$

Our recurrence relation for generating orthonormal polynomials and their derivatives ( $m = 1, 2, \dots$ ) (or their integrals with  $m = -1, -2, -3, \dots$ ) are carried out by:

$$P_{i+1}^{(m)}(E) = \gamma_{i+1} [E - \mu_{i+1}] P_i^{(m)}(E) - (1 - \delta_{i0}) \nu_i P_{i-1}^{(m)}(E) + m P_i^{(m-1)}(E), \quad (7)$$

where  $\mu_i$  and  $\nu_i$  are recurrence coefficients, and  $\gamma_i$  is a normalizing coefficient, defined by scalar products of given data. One can generate  $P_i^{(m)}(E)$  recursively. The polynomials satisfy the following orthogonality relations:

$$\sum_{i=1}^M w_i P_k^{(0)}(E_i) P_l^{(0)}(E_i) = \delta_{k,l}$$

over the discrete point set  $\{E_i, i = 1, 2, \dots, M\}$ . All the calculations for the sake of uniformity are carried out for  $E$  in  $[-1, 1]$ , i.e. after the input interval is transformed to the unit interval. We remark some advantages of OPEM: It uses unchanged the coefficients of the lower-order polynomials; it avoids the procedure of inversion of the coefficient matrix to obtain the solution. All these features shorten the computing time and assure the optimal solution by the criteria (2) and (4).

### 3.2 Usual expansion criteria

The inherited errors in usual coefficients are given by the inherited errors in orthogonal coefficients:

$$\Delta c_i = \left( \sum_{k=1}^L (c_i^{(k)})^2 \right)^{1/2} \Delta a_i, \quad (8)$$

$$\Delta a_i = \left[ \sum_{k=1}^M P_i^2(E_k) w_k (F_k - F_k^{\text{appr}})^2 \right]^{1/2}. \quad (9)$$

where coefficients  $c_i^{(k)}$  are defined by the expansion of polynomials in ordinary polynomial basis

$$P_k = \sum_{i=0}^k c_i^{(k)} E^i, k = 0, \dots, L \quad (10)$$

and explicitly constructed by recurrence relation in [12].

The procedure is iterative because of the evaluation of derivatives on every iteration step. We note that in every iteration step the algorithm find the best approximation

Table 1: **OPEM approximations results for every step approximation**

$k^{it}$	1	2	3	4	5	6	7	8	9
$L(2 \div 10)$	7	6	6	6	6	5	6	5	6
$\chi^2 * 10^{-1}$	5.61	4.24	3.99	3.79	3.77	6.81	3.75	6.65	3.63
$\max (F_a - F_c) $	14.96	3.48	6.75	4.80	4.63	0.03	6.16	0.08	9.33

using given before criteria. Having the optimal degree  $L_a$  we continue with finding the optimal  $L_c$ . We are asking for the **minimal value** of the maximal distance between functions, evaluated by orthonormal and usual expansions through all iterations.

$$\max |(F_{a,k}^{\text{appr}} - F_{c,k}^{\text{appr}}|, k = 1, \dots, M \quad (11)$$

## 4 Approximation results

The main important results with approximation degrees  $2 \div 10$  for iterations  $1 \div 9$  are presented in Table 1 for characteristics: number of iterations, number of polynomials,  $\chi^2$ , and  $\max |(F_a - F_c)|$ . We see from the Table, that for iterations  $2 \div 5$  with optimal number  $L_a = 6$  the results are good for both expansions and for 8- th iteration with optimal number  $L_c = 8$  the usual expansion is also good. We present on figures the three curves – given(B), approximated by orthogonal polynomials (C) and received from it by usual polynomials (D) at different iteration steps. Below the figures 3 and 4 show the orthogonal (C) and usual (D) approximations for 4-rd ( $L = 6$ ) and 8-th ( $L = 5$ ) iterations.




Fig. 3: Orthogonal approximation (C) and usual one (D)  $L = 6, k^{it} = 4$




Fig. 4: Orthogonal approximation (C) and usual one (D)  $L = 5, k^{it} = 8$

The Table 2 presents the approximation at 5 – th degree in 8 – th iteration of  $M = 18$  given values of following characteristics: energy  $E$ , distribution  $F$ ,  $\sigma_E$  and  $\sigma_{F(E)}$ , and from 5 – th column – the approximating values with orthonormal coefficients  $F_a^{\text{appr},5}$ , approximating values with usual coefficients  $F_c^{\text{appr},5}$ , differences  $\Delta(F_a, F_c) = (F_a^{\text{appr},5} - F_c^{\text{appr},5})$ , total variance  $S(5)$  (equation (1)). The Table 2 shows good coincidence between two descriptions. For comparison we can see the previous results for OPEM applications in [10, 11, 12].

Table 2: OPEM approximation of contact water energy data

No.	$E[ev]$	$F(E)$	$\sigma_E$	$\sigma_F$	$F_a^{appr,5}$	$F_c^{appr,5}$	$\Delta(F_a, F_c)$	S
1	0.1395	2.820	0.025	0.72	2.421	2.503	8.169-02	2.2072
2	0.1392	3.627	0.025	1.43	2.721	2.799	7.796-02	2.9469
3	0.1388	2.822	0.025	1.43	3.192	3.266	7.420-02	2.2173
4	0.1367	3.227	0.025	1.08	4.408	4.484	7.614-02	1.8114
5	0.1335	4.035	0.025	1.08	4.272	4.353	8.125-02	1.3297
6	0.1309	4.035	0.025	1.08	3.467	3.549	8.161-02	1.3126
7	0.1287	3.632	0.025	1.43	2.840	2.905	6.474-02	2.6050
8	0.1265	3.200	0.025	0.72	2.534	2.583	4.910-02	0.9395
9	0.1235	2.422	0.025	0.72	2.861	2.932	7.089-02	0.5500
10	0.1210	2.017	0.025	1.43	3.821	3.889	6.886-02	3.4402
11	0.1188	4.840	0.025	1.08	5.091	5.137	4.575-02	5.1487
12	0.1157	8.470	0.025	1.43	7.259	7.291	3.272-02	8.2753
13	0.1127	10.887	0.025	1.43	9.290	9.334	4.365-02	5.3774
14	0.1097	12.095	0.025	2.15	10.647	10.700	5.320-02	4.6238
15	0.1069	9.677	0.025	1.08	10.750	10.793	4.292-02	6.4789
16	0.1041	6.452	0.025	1.08	9.243	9.276	3.293-02	15.8508
17	0.1012	5.242	0.025	0.72	5.569	5.601	3.178-02	6.0766
18	0.0975	4.030	0.025	1.08	-2.384	-2.347	3.714-02	86.5354

## 5 Conclusions

- We have developed new version of OPEM algorithm (7) and Fortran77 package to include errors in both variables defined new “total variance” (1) according (2) and (4), and taking into account the inherited errors (8),( 9) in coefficients.
- The results show that the orthonormal and usual expansions values (5) are close to given ones in the whole interval. The optimal approximations of contact (wetting) energy show good **accuracy and stability**. We received suitable descriptions of the energy variations useful for further investigations with typical formes of approximated curves for different water probes.

## References

- [1] *A. Antonov, L. Todorova* // Compt. Rendus Acad. Bulg.Sci.1995. **V.48**. P.21-24.
- [2] *L. Todorova, A. Antonov* // Compt. Rendus Acad. Bulg.Sci. 2000. **V.53** P.43-45.
- [3] *A. Antonov, T.Galabova, L.Todorova, A.Tomov* // in Commun. Franco-Bulgare OM, 1993. **V.1**.
- [4] *D. Bonn, D. Ross* // Wetting transitions, Rep. Progr. Phys.2000. **V.64** P.1085;  
*N. A. Fuchs* // Evaporation and droplet growth in gaseaus media, Pergamon, London, 1959.
- [5] *P.R. Bevington* // Data Reduction and Error Analysis for the Physical Sciences McGraw-Hill ( New York) 1969.
- [6] *G. Jones* // Preprint TRI-PP-92-31 ,A 1992; *J. Orear* // Am. J. of Physic.s 1982. **V. 50** P. 912 ; *M.Lybanon* // Am. J. Physics. 1984. **V. 52** P. 276.
- [7] *G. Forsythe* // J. Soc. Ind. Appl.Math. 1957. **V. 5** P. 74-87.
- [8] *N. Bogdanova* // Commun. JINR Dubna, E11-98-3, 1998.
- [9] *N.Bogdanova, St.Todorov* // IJMPC, 2001. **V.12** P. 117-127.
- [10] *N. Bogdanova* // BPU6 Conf., Istanbul, August 2006, in 2007 AIP proceedings, edited by S.A.Cetin , I.Hikmet, 978-0-735400404-5/07.
- [11] *N. Bogdanova, St. Todorov* // BPU7 Conf., Alexandroupolis, Greece, Sept. 2009, in 2010 AIP proceedings, edited by Angelopoulis A, Takis Fildisis,ISBN:978-0-7354-0740-4;ISSN(print):0094-243X;ISSN(online):1551-7616.
- [12] *N.Bogdanova, St.Todorov*// MMCP 2009, Dubna, LIT, in Bulletin of PFUR, Series Mathem.Inform. Sciences.Physics. 2011. **V.3(2)** P. 63-67.